
US ERA ARCHIVE DOCUMENT

The CCRUSH Study: Characterization of Coarse and Fine Particulate Matter in Colorado

Michael Hannigan







I would like to thank the following people for their help:

Prof. Jana Milford (advisor, PI, CU)

Prof. Shelly Miller (advisor, PI, CU)

Prof. Jennifer Peel (PI, CSU)

Prof. Bill Navidi (PI, CSM)

Nicholas Clements (PhD Student, CU)

Prof. Noah Fierer (professor, CU)

Bob Bowers (post-doc, CU/UH)

John Ortega (post-doc, CU/NCAR)

Ricardo Piedrahita (post-masters, CU)

Allison Moore (undergraduate, CU)

Kelly Albano (undergraduate, CU)

Paul Monteford (graduate, CU)

Lisa Coco (undergraduate, UNC)

Dan Welsh (undergraduate, UNC)

Kelli Fischer (undergraduate, CU)

Kasey Wachtendorf (undergraduate, CU)

Berkeley Almand (PhD student, CU)

Brian Hancz (lab tech, CU)

Prof. Constantinos Sioutas (VI design, USC)

Prof. James Schauer (ECOC analysis, UW)

Bradley Rink (CDPHE)

Pat McGraw (CDPHE)

Jenny Eav (graduate student, UCB)

Bounkheana Chhung (undergraduate, CU)

Zev Ross (Zev Ross Spatial Analysis)

Tiffany Duhl (post-doc, CU)

Mingjie Xie (PhD student, CU)

Liza Boyle (PhD student, CU)

Custodial and front office staffs at the monitoring sites





Colorado Department of Public Health and Environment
















What is **coarse particulate matter (PM_{10-2.5})**?



1. Introduction

2. Methods

3. Results


4. Discussion






Why do we care about *coarse particles*?

Epidemiological studies show exposure to both PM_{2.5} and PM_{10-2.5} are detrimental to human health!






(Park and Wexler, 2008)

ALSO!
Cloud Formation
Microbe Transport
Carbon cycle
Mineral/dust cycles
Radiative Forcing





The Colorado Coarse Rural-Urban Sources and Health (CCRUSH) Study




1. Introduction 2. Methods 3. Results

4. Discussion





How did we measure particulate matter?



1. Introduction

2. Methods


3. Results

4. Discussion




Where did we measure particulate matter?









Mass Concentrations – Statistical Summary





Mass Concentrations – Temporal Trends






(* - significant difference)

1. Introduction 2. Methods 3. Results

4. Discussion



Mass Concentrations – Spatial Trends



1. Introduction


2. Methods


3. Results

4. Discussion



Mass Concentrations – Meteorological Trends





1. Introduction

2. Methods


3. Results

4. Discussion






Mass Concentrations – Meteorological Trends





Mass Concentrations – Bias Analysis

$$\begin{split} PM_{10-2.5} \ Case \ 1 &= PM_{10,FDMS} - PM_{2.5,FDMS} = PM_{10-2.5}NV + PM_{10-2.5}SV \\ PM_{10-2.5} \ Case \ 2 &= PM_{10,FDMS} - PM_{2.5} = PM_{10-2.5}NV + PM_{10}SV \\ PM_{10-2.5} \ Case \ 3 &= PM_{10} - PM_{2.5,FDMS} = PM_{10-2.5}NV - PM_{2.5}SV \\ PM_{10-2.5} \ Case \ 4 &= PM_{10} - PM_{2.5} = PM_{10-2.5}NV \end{split}$$



1. Introduction


2. Methods

3. Results

4. Discussion



Mass Concentrations - Bias Analysis



1. Introduction

2. Methods

3. Results

4. Discussion



Mass Concentrations – Bias Analysis

CPHE data is simulated by Case 3: PM₁₀(no-FDMS)-PM_{2.5}(FDMS)


 \bullet Case 3 underestimates $PM_{10-2.5}$ by the mass concentration of semi-volatile PM_{10}

Method for correcting CDPHE data:

- Model PM_{2.5} semi-volatile concentrations based on the total PM_{2.5}
- Estimate the non-volatile fraction of PM2.5 by subtracting the modeled semi-volatile values from the total PM_{2.5} concentrations
- Subtract the non-volatile PM_{2.5} from the non-volatile PM₁₀ concentrations to estimate PM_{10-2.5} with Case 4
- Case 4 underestimates by the semi-volatile fraction of PM_{10-2.5}, which is very low (0.1 µg/m³), so give a good approximation of Case 1
- Using non-volatile (or base) concentrations of TEOM instruments for estimating $PM_{10-2.5}$ seems like best low-cost option for monitoring in areas with low semi-volatile $PM_{10-2.5}$ concentrations



Mass Concentrations - Bias Analysis



Model based on 10 months of data collected from Oct 2011 to July 2012, but how do we know it should apply to 3 years of CCRUSH data?

1. Introduction 2. Methods 3. Results


5. Questions

4. Discussion



Mass Concentrations – Bias Analysis

Bootstrapped random 10 month segments of the ALS time series and performed regression between $PM_{2.5}$ and semi-volatile $PM_{2.5}$




1. Introduction 2. Methods 3. Results

4. Discussion



Mass Concentrations – Bias Analysis



1. Introduction

2. Methods


3. Results

4. Discussion



5. Questions


Elemental Concentrations - Concentrations and Enrichment



1. Introduction 2. Methods 3. Results 4. Discussion




Elemental Concentrations - Concentrations and Enrichment






Elemental Concentrations - Concentrations and Enrichment





Elemental Concentrations – Source Apportionment




1. Introduction 2. Methods 3. Results

4. Discussion



Bulk ECOC – Summary



1. Introduction


2. Methods


3. Results


4. Discussion




Bulk ECOC - Summer Peak 4 Shift














DNA Analysis - Microbe Species and Source Apportionment



1. Introduction

2. Methods


3. Results

4. Discussion





ROS Assay - Summary





CCRUSH Study Results Summary

- (1) Temporal trends are stronger for PM_{10-2.5} and PM_{2.5}, related to increased rate of removal due to sedimentation
- (2) PM_{10-2.5} and PM_{2.5} show opposite trends with wind speed and relative humidity, both fractions show the urban core of Denver as the most significant source region
- (3) Biases from using various TEOM models were shown to significantly influence total concentration and spatial statistics
- (4) Presented a method for removing biases from TEOM monitors in Denver
- (5) Elemental concentrations revealed five factors explaining different sources of PM_{10-2.5} and PM_{2.5} in Denver and Greeley: road salt, crustal emissions, coal emissions, vehicle wear emissions, and a source related to catalysts found in vehicles and refineries
- (6) OC peak fractions show a shift in dominant organic carbon type from Peak 2 to Peak 4 during the summer months at all sites
- (7) Bacterial species identified leaf surfaces and soil to be dominant sources in Denver, and cow fecal matter as the dominant source in Greeley
- (8) ROS assay results show an order of magnitude difference between $PM_{10-2.5}$ and $PM_{2.5}$ fractions, with the larger response seen for $PM_{2.5}$ samples



Future Work

- (1) Finalize ECOC data and complete analysis of monthly TSP samples
- (2) Perform endotoxin analysis
- (3) Finish analyzing samples for total microbe counts via flow cytometry
- (4) Characterize the water-soluble organic carbon fraction of collected particles
- (5) Re-analyze the three-year time series of total and semi-volatile mass concentrations

Health Effects and Characterization of Urban and Rural Coarse Particulate Matter in Northeastern Colorado

Jennifer L. Peel, PhD, MPH



EPA STAR Meeting March 18, 2013








CCRUSH - Health Effects Analysis

 Objective: Examine the association of short-term exposure coarse PM mass with respiratory and cardiovascular hospital visits in Denver and Greeley, CO



CCRUSH: Methods - Denver


- Daily non-elective hospital admissions, emergency department visits for all ages, 2009-2011
- 20 miles radius around the city Center





CCRUSH: Methods - Greeley

- Daily non-elective hospital admissions, emergency department visits for all ages, 2009-2011
- 20 miles radius around the city Center





CCRUSH: Methods

- Cardiovascular, Respiratory
- COPD (ICD-9 490-492, 496)
- Asthma (493)
- Pneumonia (480-486)
- Upper Respiratory Infection (460-466, 477)
- Ischemic heart disease (410-414)

CCRUSH: Health models

- Poisson generalized additive models (GAMs)
 - Adjust for long term temporal trends (12 df per year), day of week, federal holidays, temperature (3 df), dewpoint (3 df)
- Lag 0 for CVD
- Lag 0, Distributed lag (0-1, 0-4) for respiratory outcomes
- RRs and 95% Cis per IQR increase
 - 9.9μg/m³ for Denver
 - 9.0µg/m³ for Greeley





CCRUSH: Preliminary Results

Results primarily null

Low daily counts, limited power, wide confidence intervals

 Suggestion of stronger effects for asthma, particularly for Greeley

CCRUSH: Ongoing Work

- Finalizing ED visit data for Denver
- Additional pollutants: PM₁₀, PM_{2.5}, ozone, CO, NO₂
- Alternative lag structures
- Additional health outcomes: adverse birth outcomes (preterm birth, low birth weight)
- Sensitivity analyses: alternate control for temporal trends, meteorology; case-crossover analyses



Thank you!



References

Brunekreef and Forsberg. European Respiratory Journal, 26: 309-318 (2005).

Clements et al. Aerosol Science and Technology, 46(1): 108-123 (2012).

Despres et al. Tellus Sereis B (2012).

Harrison et al. Environmental Science and Technology, 46: 6523-6529 (2012).

Huang et al. Atmospheric Environment, 66: 17-24 (2013).

Park and Wexler. Aerosol Science, 39: 266-276 (2008).